

AperTO - Archivio Istituzionale Open Access dell'Università di Torino

Correlates of violent suicide attempts in patients with bipolar disorder

This is the author's manuscript

Original Citation:

Availability:

This version is available <http://hdl.handle.net/2318/1717398> since 2019-11-24T17:11:33Z

Published version:

DOI:10.1016/j.comppsy.2019.152136

Terms of use:

Open Access

Anyone can freely access the full text of works made available as "Open Access". Works made available under a Creative Commons license can be used according to the terms and conditions of said license. Use of all other works requires consent of the right holder (author or publisher) if not exempted from copyright protection by the applicable law.

(Article begins on next page)

Journal Pre-proof

Correlates of violent suicide attempts in patients with bipolar disorder

Gianluca Rosso, Umberto Albert, Stefano Bramante, Elena Aragno,
Federica Quarato, Gabriele Di Salvo, Giuseppe Maina



PII: S0010-440X(19)30059-8

DOI: <https://doi.org/10.1016/j.comppsy.2019.152136>

Reference: YCOMP 152136

To appear in:

Please cite this article as: Rosso G, Albert U, Bramante S, Aragno E, Quarato F, Di Salvo G, Maina G, Correlates of violent suicide attempts in patients with bipolar disorder, *Comprehensive Psychiatry* (2019), doi: <https://doi.org/10.1016/j.comppsy.2019.152136>

This is a PDF file of an article that has undergone enhancements after acceptance, such as the addition of a cover page and metadata, and formatting for readability, but it is not yet the definitive version of record. This version will undergo additional copyediting, typesetting and review before it is published in its final form, but we are providing this version to give early visibility of the article. Please note that, during the production process, errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

© 2019 Published by Elsevier.

Title: Correlates of violent suicide attempts in patients with bipolar disorder.

Author names and affiliations: Gianluca Rosso^a, Umberto Albert^c, Stefano Bramante^{a,b}, Elena Aragno^{a,b}, Federica Quarato^{a,b}, Gabriele Di Salvo^{a,b}, Giuseppe Maina^{a,b*}

- a. Psychiatric Unit, San Luigi Gonzaga Hospital, Regione Gonzole 10, 10043, Orbassano, Turin, Italy.
- b. Department of Neurosciences “Rita Levi Montalcini”, University of Turin, Via Cherasco 15, 10126, Turin, Italy.
- c. Department of Medicine, Surgery and Health Sciences, Psychiatric Section, University of Trieste, Via Guglielmo de Pastrovic 4 -34128, Trieste, Italy.

***Corresponding author**

Giuseppe Maina

Psychiatric Unit, San Luigi Gonzaga Hospital, Regione Gonzole 10, 10043, Orbassano, Turin, Italy.

Telephone number: +39 011 9026517

Fax number: +39 011 9026669

E-mail address: giuseppe.maina@unito.it

Highlights:

- Cross-sectional study to analyze violent suicide attempts in bipolar disorder;
- 30.6% of attempters used a violent suicide method;
- More men in the group of violent suicide attempters;
- Association between obesity and violent suicide.

Abstract

BACKGROUND: Suicide is one of the leading causes of death in bipolar disorder (BD); violent suicide attempts are associated with the highest level of lethality. We aimed to evaluate factors related to the risk of violent

suicide in a large naturalistic sample of patients with BD; in addition, we analyzed the rates of lifetime suicide attempts and the variables associated with suicidal behavior.

METHODS: We recruited 847 patients with BD. Patients were grouped according to whether they had a lifetime history of suicide attempts and, among suicide attempters, subjects who had used a violent suicide method were compared with those who had attempted suicide with a nonviolent method. Comparisons were performed using χ^2 tests for categorical variables and ANOVA for continuous variables. Logistic regression (LogReg) was used to identify explanatory variables associated with violent suicide attempts (dependent variable).

RESULTS: Two hundred and two patients (24%) had a lifetime history of suicide attempts. Subjects with at least one lifetime suicide attempt showed longer duration of illness (22.4 ± 14.1 years vs 19.9 ± 14.2 years: $p = 0.028$), more lifetime hypomanic episodes (3.3 ± 4.3 vs 2.3 ± 3.1 : $p = 0.001$), more lifetime depressive episodes (6.0 ± 4.4 vs 4.7 ± 4.1 : $p < 0.001$), higher rates of lifetime psychiatric comorbidities (50.0% vs 41.3%: $p = 0.029$), higher rates of lifetime medical comorbidities (58.0% vs 48.9%: $p = 0.028$) and higher rates of reduced HDL cholesterol (46.2% vs 36.7%: $p = 0.030$). Among suicide attempters, fifty-two patients (30.6%) attempted suicide with a violent method. We found more men in the group of violent suicide attempters than in the group of nonviolent suicide attempters (65% vs 28%; $p < 0.001$). Moreover subjects with previous violent attempts showed higher mean values of weight (80.5 ± 18.3 vs 69.4 ± 14.7 : $p < 0.001$), body mass index (27.8 ± 5.6 vs 25.2 ± 4.7 : $p < 0.003$) and waist circumference (98.7 ± 18.5 vs 92.4 ± 14.3 : $p = 0.032$). The LogReg analysis confirmed the association of violent attempts with male gender ($p < 0.001$; Phi: 0.35) and higher waist circumference ($p < 0.001$; Cohen's d : 0.39).

LIMITATIONS: In our research we analyzed lifetime suicide attempts, but the sample does not include completed suicides, meaning that we are unable to test whether the results are generalizable to suicide deaths. Moreover, some relevant variables, such as medical comorbidities/metabolic parameters at the time of suicide attempts and previous medication, were not collected. Another limitation concerns the heterogeneity of recruited patients in terms of clinical characteristics (e.g.: medical conditions, drug treatments), with potential confounding factors.

CONCLUSIONS: The present study confirms the association between male gender and violent suicide and suggests a correlation between obesity and the use of violent suicide methods. The relationship between obesity and suicidal behaviour is worthy of interest and deserves to be explored by further studies.

Keywords

Bipolar disorder; suicide attempts; violent suicide.

Manuscript

1. Introduction

Bipolar disorder (BD) is a recurrent and chronic disease characterized by the occurrence of manic (or hypomanic), depressive, or mixed episodes. According to the World Health Organization, BD is one of the world's ten most disabling conditions [1, WHO 2008]. Suicide is one of the leading causes of death in BD: as reported by the most recent guidelines [2, Yatham et al., 2018], approximately 6%-7% of identified patients with BD dies by suicide, 43% report suicidal ideation, 21% suicide plan and 16% suicide attempt within the past year. Existing studies suggest that 13.2%-17.5% of patients with BD with a lifetime history of suicide attempt performs them with violent methods, such as hanging, shooting or jumping from heights; these subjects, despite representing a minority, need to be precociously recognised, as these methods are associated with the highest level of lethality [3, Cassidy, 2011].

An extensive literature has deepened socio-demographic, clinical, environmental, genetic and neurobiological factors related to violent suicide methods in psychiatric disorders [4, Ludwig et al., 2018]. It has been well established that men use violent methods more often than women, both in suicide and suicide attempts [5, Bradvik, 2007] [6, Mergl et al., 2015]. Furthermore, higher impulsivity and aggression levels as well as substance abuse disorder have been associated with violent suicide [7, Zalsman et al., 2006] [8, Ilgen et al., 2010] [9, Zupanc et al., 2013] [10, Sheehan et al., 2015]. A few studies investigated the correlation between violent suicide and environmental factors, including seasonality, temperature, relative humidity, rainfall, hours of sunshine and pressure. A significant positive correlation was found between violent attempts and ambient temperature and number of sunshine hours [11, Lin et al., 2008] [12, Vyssoki et al., 2014]. Recent research works deepened the relation between violent suicide and serotonin axis, including candidate gene studies and gene expression studies. In a candidate gene study with a sample of schizophrenic patients, a significant association was found between violent suicide and the genotype frequencies of promoter of the serotonin-transporter gene (5HTTLPR) polymorphism [13, Balye et al., 2003]. Other studies evaluated differences in the promoter polymorphism of the tryptophane hydroxylase (TPH) between violent and nonviolent suicide attempters, with conflicting results [Stefulj et al., 2005] [Stefulj et al., 2006]. In order to identify neurobiological factors related to violent suicide, the implication of cytokines in the pathogenesis of suicide has been recently investigated. Interleukin-6, measured in cerebrospinal fluid (CSF), peripheral blood, and postmortem brain tissue, has shown robust associations with violent suicide and suicide completion [16, Gananca et al., 2016]. Furthermore, the relationships between metabolic parameters and violent suicide have been extensively studied. A study conducted by Tanskanen and colleagues demonstrated that serum total cholesterol concentration was positively related to violent but not nonviolent suicide; the same trend could be shown for the concentration of high density lipoprotein cholesterol, but not reaching statistical significance [17, Tanskanen et al., 2000]. Conversely, two studies performed by Marcinko and colleagues found significantly lower cholesterol serum levels in violent suicide attempters compared with suicide attempters using nonviolent methods [18, Marcinko et al., 2004] [19, Marcinko et al., 2005]. A similar case-control study looked at women with a history of suicide attempts and compared the serum cholesterol concentration between violent suicide attempters and nonviolent suicide attempters: again, a lower concentration was predictive of violent methods within this sample of patients [20, Vevera et al., 2003]. Two postmortem studies, relevant to the question of fatty acid composition and violent suicide, reported the following findings: significantly lower cholesterol levels in the prefrontal cortex of violent suicide attempters, significantly higher cholesteryl ester hydrolase (LIPA) expression in violent suicide attempters, a significantly lower cholesterol/phospholipid level ratio in violent suicide attempters and increased levels of phospholipids in violent suicide attempters when compared with nonviolent suicide attempters [21, Freemantle et al., 2013] [22, Lalovic et al., 2007].

Risk factors for violent suicide, with special reference to BD, were investigated by few studies, mainly conducted with small samples and heterogeneous methods. Available data suggest that male gender [23, Perroud et al., 2007] [24, Nivoli et al., 2011] [25, Dell'Osso et al., 2017], onset of BD with a hypo/manic episode [26, Neves et al., 2008] [27, Schaffer et al., 2016], S allele of the serotonin transporter gene (5-HTTLPR) polymorphisms [26 Neves et al., 2008] and single nucleotide polymorphisms of the CLOCK and TIMELESS genes [28, Pawlak et al., 2015] seem to be correlated with an increased risk of suicide with violent methods. A study performed by our research group analyzed the correlation between metabolic syndrome, lipid profile and suicide with violent methods in bipolar patients, but no significant differences were found in violent suicide attempters compared to nonviolent ones [29, D'Ambrosio et al., 2012].

The primary goal of this study was to evaluate in a large naturalistic sample of patients with BD potential factors related to the risk of suicide with violent methods. In addition, we sought to examine in our sample the rates of lifetime suicide attempts and the variables associated with suicidal behavior.

2. Material and methods

2.1 Study design and patients

Data derive from an independent cross-sectional observational study aimed to analyze course characteristics, medical conditions and response to treatments in patients with BD.

Subjects for this study were recruited from all patients with a principal diagnosis of BD type I, II or Not Otherwise Specified (NOS) (DSM-IV-TR, DSM-5) [30, American Psychiatric Association 2000] [31, American Psychiatric Association 2013] consecutively admitted to the Department of Neuroscience, University of Turin (Italy), from January 2006 to February 2019.

Potential participants were thoroughly explained aims and study procedures and had to give their written consent before participation; exclusion criteria included age <18 and refusal to consent participating in the study. The protocol was approved by the local Ethical Committees.

2.2 Assessments and procedures

All diagnoses were confirmed by means of the Mini-International Neuropsychiatric Interview (MINI) [32, Sheehan et al., 1998]. At the study entry, general socio-demographic information and clinical data were collected for each subject. Clinical characteristics such as age at onset, duration of illness, number of previous manic/depressive episodes, psychiatric comorbidity, psychiatric family history, current medication and psychopharmacological treatments were ascertained either from clinical charts or by direct questioning the study participants. History of suicide attempt, defined as a self-destructive behavior with the intention of ending one's life, independently of the resulting damage, [33, O'Carroll et al., 1996] was retrospectively assessed for each patient, focusing on the modality of suicidal behavior. According to Stenbacka et al. [34, 2015], the suicide attempt method was defined violent (hanging, shooting, jumping from heights, moving train, cutting, drowning) or nonviolent (poisoning). Individuals who had made more than one attempt were classified according to the violence of their most violent attempt.

All subjects received a medical examination; a specific evaluation of metabolic parameters has been conducted for each patient: weight was measured fasting and undressed, height barefoot, waist circumference at midway between the inferior margin of the ribs and the superior border of the iliac crest, at minimal inspiration. Body mass index (BMI) was calculated as the body weight in kilograms divided by the square of the body height in metres. Two blood pressure measurements were obtained by using a mercury sphygmomanometer: the first with the subject in a lying position, the second with the subject in a seated position at least two minutes after the first measurement. The mean blood pressure of the two measurements was used. A blood sample (including cholesterol, glucose, triglycerides and HDL-c) was drawn in the morning (7:00 am), when patients were fasting for previous 10 hours. Metabolic syndrome has been evaluated using definition of the International Diabetes Federation (IDF) Task Force on Epidemiology and Prevention, American Heart Association and National Heart, Lung and Blood Institute [35, Grundy et al., 2005]: metabolic syndrome was present if three or more of the following five criteria were met:

- Abdominal obesity: waist circumference ≥ 94 cm in men and ≥ 80 cm in women;
- High blood pressure: systolic pressure ≥ 130 mmHg and/or diastolic pressure ≥ 85 mmHg or on antihypertensive medication;
- High fasting glucose: ≥ 100 mg/dl or on glucose-lowering medication;
- Hypertriglyceridemia: ≥ 150 mg/dl or on lipid-lowering medication;

- Low HDL-C: < 40 mg/dl in men and < 50 mg/dl in women.

2.3 Statistical analysis

Socio-demographic and clinical features of the patients were summarized as mean and standard deviation (SD) for continuous variables and frequency and percentage for categorical variables. Patients were grouped according to whether they had a lifetime history of suicide attempts or they had never attempted suicide in their life. Further, patients who had used a violent suicide method were compared with those who had attempted suicide with a nonviolent method.

The normality of data distribution was evaluated by using Shapiro-Wilk and Kolmogorov-Smirnov tests. Comparisons were performed using χ^2 tests for categorical variables and ANOVA for continuous variables. Furthermore, logistic regression (LogReg) was used to identify explanatory variables associated with violent suicide attempts (dependent variable).

The results from every statistical comparison of the treatment groups were presented as 2-sides p values rounded to 3 decimal places. The criterion for statistical significance in all comparison was a p value < 0.05. All statistical analyses were performed by SPSS software version 22.0.

3. Results

Eight-hundred fifty-three patients with BD were asked to participate; six refused their consent. Among the 847 patients recruited, 5 (0.6%) were excluded from the research due to lack of data about lifetime suicide attempts.

Ultimately, we completed the analysis using 842 subjects. The demographic and clinical characteristics of the total sample are given in Table 1. The sample is representative for the population of patients with BD: 58.7% of the patients were females, the majority of the sample (54.5%) had bipolar II disorder, the mean age at onset of BD was 29.3 ± 12.4 years, the mean duration of illness was 20.5 ± 14.2 years. Two hundred and two patients (24%) had a lifetime history of suicide attempts; among suicide attempters, 22 subjects (10.9%) had made more than one attempt in their life. The demographic and clinical differences between suicide attempters and non attempters are summarized in Table 1: subjects with at least one lifetime suicide attempt showed longer duration of illness (22.4 ± 14.1 years vs 19.9 ± 14.2 years: p 0.028), more lifetime hypomanic episodes (3.3 ± 4.3 vs 2.3 ± 3.1 : p 0.001), more lifetime depressive episodes (6.0 ± 4.4 vs 4.7 ± 4.1 : p < 0.001), higher rates of lifetime psychiatric comorbidities (50.0% vs 41.3%: p 0.029), higher rates of lifetime medical comorbidities (58.0% vs 48.9%: p 0.028) and higher rates of reduced HDL cholesterol (46.2% vs 36.7%: p 0.030).

The suicide attempters' group was further analyzed according to whether they had employed a violent or a nonviolent suicide attempt; within the attempters' group 32 subjects (15.8%) were excluded due to lack of data about the modality of suicidal behavior. The results are shown in Table 2. Fifty-two patients (30.6%) attempted suicide with a violent method. Violent suicidal attempts consisted respectively in jumping from heights (n=17), hanging (n=13), abdominal knife wounds (n=7), gunshot wounds (n=4), gas poisoning (n=4), car accident (n=3), others (n=4). In our sample, nonviolent method was drug overdose (n=118). There was a statistically significant difference in sex distribution (p: <0.001), with more men in the group of violent suicide attempters than in the group of nonviolent suicide attempters (65% vs 28%; p: <0.001). Moreover, we found significant differences within metabolic parameters: subjects with previous violent attempts, in comparison to patients with nonviolent attempts, showed higher mean values of weight (80.5 ± 18.3 vs 69.4 ± 14.7 : p <0.001), BMI (27.8 ± 5.6 vs 25.2 ± 4.7 : p <0.003) and waist circumference (98.7 ± 18.5 vs 92.4 ± 14.3 : p 0.032).

There were no further significant differences within other socio-demographic and clinical characteristics between the two groups. The LogReg analysis confirmed the association of violent attempts with male gender ($p: <0.001$; Phi: 0.35) and with higher waist circumference ($p: <0.001$; Cohen's $d: 0.39$). The results of the LogReg analysis are described in Table 3; independent variables included were gender, BMI, weight, abdominal circumference. The LogReg model was statistically significant, $\chi^2(4) = 29.590$, $p < .001$. The model explained 28.0% (Nagelkerke R^2) of the variance in response and correctly classified 71.6% of cases. Since the LogReg ruled out weight and BMI as independent risk factors, monovariate analyses have been conducted in gender subgroups. In male subgroup no differences emerged between violent and nonviolent suicide attempters; in female subgroup there are statistically significant differences between violent and nonviolent suicide attempters in terms of higher mean body weight ($73,2 \text{ kg} \pm 14,5$ vs $65,65 \pm 12,9$; $p=.034$) and BMI ($27,7 \pm 4,3$ vs $25,1 \pm 4,9$; $p=.047$).

4. Discussion

The main objective of this study was to analyze socio-demographic and clinical factors related to violent suicide attempts in a large naturalistic sample of patients with BD; in addition, we analyzed the rates of lifetime suicide attempts and the variables associated with suicidal behavior.

The prevalence of lifetime suicide attempts found in our sample (24%) is consistent with those reported by previous studies [36, Tondo et al., 2003] [37, Manchia et al., 2013] [27, Schaffer et al., 2015]. Some clinical variables (longer duration of illness, more lifetime hypomanic and depressive episodes, higher rates of lifetime psychiatric and medical comorbidities) were showed to be associated with greater likelihood of lifetime suicide attempts: these results were expected and confirm the most important studies recently carried out on this topic [38, Tidemalm et al., 2014] [27, Schaffer et al., 2015]. Furthermore, a correlation emerged between reduced HDL cholesterol and lifetime suicide attempts: to date, two studies reported lower serum total cholesterol levels in BD patients with history of suicide attempt [38, Tidemalm et al., 2014] [39, Vuksan-Cusa et al., 2009], while no association was identified in a previous Italian study [29, D'Ambrosio et al., 2012]. Therefore, our finding is noteworthy and deserve further investigation in future studies.

In our sample about one-third of patients (30.6%) with history of suicidal behavior used violent methods, while previous studies showed lower rates of violent suicide attempts in BD (13.2%-17.5%) [40, Yen et al., 2008] [41, De Abreu et al., 2012] [29, D'Ambrosio et al., 2012]. There are possible explanations for the lack of correspondence between our results and available literature data. Firstly, the sample consists of patients with a high severity of illness, as our Department is a tertiary referral center located within the University General Hospital. Furthermore, investigations on suicidal behaviors in BD, including the present study, are heterogeneous, in terms of characteristics of the samples (e.g.: medications, psychiatric comorbidities), instruments used to assess suicidality and definition of violent or nonviolent methods. Comparing the subjects according to whether they had employed a violent or a nonviolent suicide method, we found significant differences across gender, with more men in the group of violent suicide attempters. Our result is consistent with previous studies that have demonstrated that the main differences in suicidal behavior between men and women is the method chosen to attempt suicide [23, Perroud et al., 2007] [24, Nivoli et al., 2011] [29, D'Ambrosio et al., 2012]. No further differences regarding socio-demographic and psychopathological features were found between the two groups. Concerning medical comorbidities and metabolic disorders, our results showed higher mean values of weight, BMI and waist circumference in the group of violent suicide attempters, while no significant difference was found in terms of serum lipid levels, glycemia, blood pressure and metabolic syndrome. The LogReg analysis confirmed also abdominal circumference as independent variable associated with violent suicide attempts; results from monovariate analyses conducted in gender subgroups showed in female group higher mean body weight and BMI values in violent attempters. To our knowledge only one previous study, performed by our research group, with a

smaller sample size, specifically evaluated the correlation between violent suicide attempts and metabolic parameters in BD: the results did not show any significant difference between violent and nonviolent suicide attempters. [29, D'Ambrosio et al., 2012]. The methodological limitations of the present study, discussed in the following paragraph, make it difficult to interpret the association between higher BMI/abdominal obesity and violent suicide methods: data on metabolic parameters at the time of suicide attempts and previous pharmacological treatments would be needed to distinguish whether BMI and waist circumference may directly influence the use of violent methods or whether subjects with history of violent attempts are more likely to be prescribed medication that leads to weight gain and abdominal obesity (e.g.: second generation antipsychotics). The hypothesis of a causal relationship between obesity and violent suicide methods could be explained by the association between obesity and impulsivity. Indeed, impulsive personality traits are prevalent characteristics in obese BD patients and some studies suggest that obesity and impulsivity together may lead to severe course of illness and worse prognosis [42, Galvez et al., 2014]. Therefore, the relationship between obesity and suicidal behaviour is worthy of interest and deserves to be explored by further methodologically rigorous studies.

Our study presents several limitations, mostly due to the cross-sectional design and to the fact that data derives from a study aimed to investigate clinical characteristics of BD and not suicidal behavior specifically. First, in our research we analyzed lifetime suicide attempts, but the sample does not include completed suicides, meaning that we are unable to test whether the results are generalizable to suicide deaths; then suicidal intent to die is difficult to ascertain retrospectively and information could be biased. Moreover, some patients have missing data about the modality of suicidal attempts and some relevant variables, such as medical comorbidities and metabolic parameters at the time of suicide attempts. Furthermore, in our analysis the impact of pharmacological treatment on suicidal behavior and on metabolic parameters was not evaluated, due to the lack of data on medications prior to study entry. Another limitation concerns the heterogeneity of recruited patients in terms of clinical characteristics (e.g.: medical conditions, drug treatments), with potential confounding factors.

Instead, the large sample size is a strength of the study. Furthermore, subjects enrolled for this study were representative of “real-world” in and outpatients with BD: this should be considered a point of strength in terms of generalizability and external validity.

5. Conclusions

In conclusion, the present study confirms the association between male gender and violent suicide; moreover, our findings suggest a correlation between obesity and the use of violent suicide methods, especially in female patients, but further research is awaited to explore and clarify this relationship. Considering that suicide is regrettably frequent in patients with BD future studies on this issue could help detecting individuals at risk of suicide attempt and take appropriate actions as early as possible.

Acknowledgements

We would like to acknowledge the staff of our Department, which helped to collect all patients' data.

Declaration on interests

Gianluca Rosso is /has been a speaker and/or has received research grants from Angelini, Janssen, Lundbeck, Otsuka.

Umberto Albert is /has been a speaker and/or has received research grants from Angelini, Innova Pharma, Neuraxpharm, Janssen, Lundbeck.

Stefano Bramante has been a speaker and has received research grants from Angelini.

Gabriele Di Salvo has been a speaker and has received research grants from Lundbeck.

Giuseppe Maina is /has been a consultant and/or a speaker and/or has received research grants from Angelini, Boehringer Ingelheim, FB-Health, Janssen, Lundbeck, Otsuka.

References

1. World Health Organization. The Global Burden of disease: 2008 update. WHO Library 2008.
2. Yatham LN, Kennedy SH, Parijth SV, Schaffer A, Bond DJ, Frey BN, Sharma V, Goldstein BI, Rej S, Beaulieu S, Alda M, MacQueen G, Milev RV, Ravindran A, O'Donovan C, McIntosh D, Lam RW, Vazquez G, Kapczynski F, McIntyre RS, Kozicky J, Kanba S, Lafer B, Suppes T, Calabrese JR, Vieta E, Malhi G, Post RM, Berk M. Canadian Network for Mood and Anxiety Treatments (CANMAT) and International Society for Bipolar Disorders (ISBD) 2018 guidelines for the management of patients with bipolar disorder. *Bipolar Disord* 2018; 20:97–170. <https://doi.org/10.1111/bdi.12609>.
3. Cassidy F. Risk factors of attempted suicide in bipolar disorder. *Suicide Life Threat Behav* 2011; 41(1):6–11. <https://doi.org/10.1111/j.1943-278X.2010.00007.x>.
4. Ludwig B, Dwivedy Y. The concept of violent suicide, its underlying trait and neurobiology: A critical perspective. *Eur Neuropsychopharmacol* 2018; 28(2):243–251. <https://doi-org.bibliopass.unito.it/10.1016/j.euroneuro.2017.12.001>
5. Brådvik L. Violent and nonviolent methods of suicide: different patterns may be found in men and women with severe depression. *Arch Suicide Res* 2007; 11(3):255–64. <https://doi-org.bibliopass.unito.it/10.1080/13811110701402611>
6. Mergl R, Koburger N, Heinrichs K, Székely A, Dittmann TM, Coyne J, Quintão S, Arensman E, Coffey C, Maxwell M, Värnik A, Audenhove C, McDaid D, Sarchiapone M, Schmidtke A, Genz A, Gusmão R, Hegerl U. What are reasons for the large gender differences in the lethality of suicidal acts? An Epidemiological Analysis in Four European Countries. *PLoS One* 2015; 10(7):e0129062. <https://doi-org.bibliopass.unito.it/10.1371/journal.pone.0129062>
7. Zalsman G, Braun M, Arendt M, Grunebaum MF, Sher L, Burke AK, Brent DA, Chaudhury SR, Mann JJ, Oquendo MA. A comparison of the medical lethality of suicide attempts in bipolar and major depressive disorders. *Bipolar Disord* 2006; 8:558–565. <https://doi-org.bibliopass.unito.it/10.1111/j.1399-5618.2006.00381.x>
8. Ilgen MA, Conner KR, Valenstein M, Austin K, Blow FC. Violent and nonviolent suicide in veterans with substance-use disorders. *J Stud Alcohol Drugs* 2010; 71 (473–479). <https://doi.org/10.15288/jsad.2010.71.473>
9. Zupanc T, Agius M, Paska AV, Pregelj P. Blood alcohol concentration of suicide victims by partial hanging. *J Forensic Leg Med* 2013; 20:(976–979). <https://doi-org.bibliopass.unito.it/10.1016/j.jflm.2013.09.011>
10. Sheehan CM, Rogers RG, Boardman JD. Postmortem presence of drugs and method of violent suicide. *J Drug Issues* 2015; 45(3):249–262. <https://dx-doi-org.bibliopass.unito.it/10.1177%2F0022042615580988>
11. Lin HC, Chen CS, Xirasagar S, Lee HC. Seasonality and climatic associations with violent and nonviolent suicide: a population-based study. *Neuropsychobiology* 2008; 57(32–37).
12. Vyssoki B, Kapusta ND, Praschak-Rieder N, Dorffner G, Willeit M. Direct effect of sunshine on suicide. *JAMA Psychiatry* 2014; 71(11):1231–1237. <https://doi:10.1001/jamapsychiatry.2014.1198>
13. Bayle FJ, Leroy S, Gourion D, Millet B, Olié JP, Poirier MF, Krebs MO. 5HTTLPR polymorphism in schizophrenic patients: further support for association with violent suicide attempts. *Am J Med Genet B Neuropsychiatr Genet* 2003; 15:119B(1):13–7. <https://doi-org.bibliopass.unito.it/10.1002/ajmg.b.10037>
14. Stefulj J, Kubat M, Bališa M, Jernej B. TPH gene polymorphism and aging: indication of combined effect on the predisposition to violent suicide. *Am J Med Genet B Neuropsychiatr Genet* 2006; 141B:139–141. <https://doi-org.bibliopass.unito.it/10.1002/ajmg.b.30271>
15. Stefulj J, Kubat M, Bališa M, Skavic J, Jernej B. Variability of the tryptophan hydroxylase gene: study in victims of violent suicide. *Psychiatry Res* 2005; 134:67–73. <https://doi-org.bibliopass.unito.it/10.1016/j.psychres.2004.04.011>
16. Gananca L, Oquendo MA, Tyrka AR, Cisneros-Trujillo S, Mann JJ, Sublette ME. The role of cytokines in the pathophysiology of suicidal behavior. *Psychoneuroendocrinology* 2016; 63:296–310. <https://doi-org.bibliopass.unito.it/10.1016/j.psyneuen.2015.10.008>
17. Tanskanen A, Vartiainen E, Tuomilehto J, Viinamäki H, Lehtonen J, Puska P. High serum cholesterol and risk of suicide. *Am J Psychiatry* 2000; 157:648–650. <https://doi-org.bibliopass.unito.it/10.1176/appi.ajp.157.4.648>

18. Marcinko D, Martinac M, Karlovic D, Loncar C. Cholesterol serum levels in violent and non-violent young male schizophrenic suicide attempters. *Psychiatr Danub* 2004; 16:161–164.
19. Marcinko D, Martinac M, Karlovic D, Filipcic I, Loncar C, Pivac N, Jakovljevic M. Are there differences in serum cholesterol and cortisol concentrations between violent and non-violent schizophrenic male suicide attempters? *Coll Antropol* 2005; 29:153–157.
20. Vevera J, Zukov I, Morcinek T, Papezova H. Cholesterol concentrations in violent and non-violent women suicide attempters. *Eur Psychiatry* 2003; 18:23–27. [https://doi-org.bibliopass.unito.it/10.1016/S0924-9338\(02\)00011-1](https://doi-org.bibliopass.unito.it/10.1016/S0924-9338(02)00011-1)
21. Freemantle E, Mechawar N, Turecki G. Cholesterol and phospholipids in frontal cortex and synaptosomes of suicide completers: relationship with endosomal lipid trafficking genes. *J Psychiatr Res* 2013; 47:272–279. <https://doi-org.bibliopass.unito.it/10.1016/j.jpsychires.2012.10.019>
22. Lalovic A, Levy E, Luheshi G, Canetti L, Grenier E, Sequeira A, Turecki G. Cholesterol content in brains of suicide completers. *Int J Neuropsychopharmacol* 2007; 10:159–166. <https://doi-org.bibliopass.unito.it/10.1017/S1461145706006663>
23. Perroud N, Baud P, Preisig M, Etain B, Bellivier F, Favre S, Reber N, Ferrero F, Leboyer M, Malafosse A. Social phobia is associated with suicide attempt history in bipolar inpatients. *Bipolar Disord* 2007; 9(7):713–21. <https://doi.org/10.1111/j.1399-5618.2007.00471.x>.
24. Nivoli AM, Pacchiarotti I, Rosa AR, Popovic D, Murru A, Valenti M, Bonnin CM, Grande I, Sanchez-Moreno J, Vieta E, Colom F. Gender differences in a cohort study of 604 bipolar patients: the role of predominant polarity. *J. Affect Disord* 2011; 133(3):443–9. <https://doi.org/10.1016/j.jad.2011.04.055>.
25. Dell’Osso B, Vismara M, Dobra C, Cremaschi L, Grancini B, Arici C, Benatti B, Buoli M, Ketter TA, Altamura AC. Clinical characterization of italian suicide attempters with bipolar disorder. *CNS Spectr* 2018; 23(4):271–277. <https://doi.org/10.1017/S1092852917000384>.
26. Neves FS, Malloy-Diniz LF, Barbosa IG, Brasil PM, Corrêa H. Bipolar disorder first episode and suicidal behavior: are there differences according to type of suicide attempt? *Braz J Psychiatry* 2009; 31(2):114–8. <http://dx.doi.org/10.1590/S1516-44462009000200006>.
27. Schaffer A, Isometsä ET, Azorin JM, Cassidy F, Goldstein T, Rihmer Z, Sinyor M, Tondo L, Moreno DH, Turecki G, Reis C, Kessing LV, Ha K, Wizman A, Beautrais A, Chou YH, Diazgranados N, Levitt AJ, Zarate CA Jr, Yatham L. A review of factors associated with greater likelihood of suicide attempts and suicide deaths in bipolar disorder: Part II of a report of the International Society for Bipolar Disorders Task Force on Suicide in Bipolar Disorder. *Aust NZJ Psychiatry* 2015; 49(11):1006–20. <https://doi.org/10.1177%2F0004867415594428>.
28. Pawlak J, Dmierzak-Weglaz M, Maciukiewicz M, Wilkosc M, Leszczynska-Rodziewicz A, Zaremba D, Kapelski P, Hauser J. Suicidal behavior in the context of disrupted rhythmicity in bipolar disorder - Data from an association study of suicide attempts with clock genes. *Psychiatry Research* 2015; 226:517–520. <https://doi-org.bibliopass.unito.it/10.1016/j.psychres.2015.01.010>
29. D’Ambrosio V, Salvi V, Bogetto F, Maina G. Serum lipids, metabolic syndrome and lifetime suicide attempts in patients with bipolar disorder. *Prog Neuropsychopharmacol Biol Psychiatry* 2012; 37(1):136–40. <https://doi.org/10.1016/j.pnpbp.2011.12.009>.
30. American Psychiatric Association. Diagnostic and Statistical Manual of Mental Disorders: DSM-IV-TR. APA 2000. <https://doi.org/10.1176/appi.books.9780890423349.5847>.
31. American Psychiatric Association. Diagnostic and Statistical Manual of Mental Disorders: DSM-V. APA 2013. <https://doi.org/10.1176/appi.books.9780890425596.744053>.
32. Sheehan DV, Lecrubier Y, Sheehan KH, Amorim P, Janavs J, Weiller E, Hergueta T, Baker R, Dunbar GC. The Mini-International Neuropsychiatric Interview (M.I.N.I.): the development and validation of a structured diagnostic psychiatric interview for DSM-IV and ICD-10. *J Clin Psychiatry* 1998; 59, 22–33. [https://doi.org/10.1016/S0924-9338\(99\)80239-9](https://doi.org/10.1016/S0924-9338(99)80239-9).
33. O’Carroll PW, Berman AL, Maris RW, Moscicki EK, Tanney BL, Silverman MM. Beyond the tower of babel: a nomenclature for suicidology. *Suicide Life Threat Behav* 1996; 26(3):237–52. <https://doi.org/10.1111/j.1943-278X.1996.tb00609.x>.

34. Stenbacka M, Jokinen J. Violent and non-violent methods of attempted and completed suicide in swedish young men: the role of early risk factors. *BMC Psychiatry* 2015; 15:196. <https://doi-org.bibliopass.unito.it/10.1186%2Fs12888-015-0570-2>
35. Grundy SM, Cleeman JI, Daniels SR, Donato KA, Eckel RH, Franklin BA, Gordon DJ, Krauss RM, Savage PJ, Smith SC Jr, Spertus JA, Costa F. American Heart Association. National Heart, Lung, and Blood Institute. Diagnosis and management of the metabolic syndrome: an American Heart Association/National Heart, Lung, and Blood Institute Scientific Statement. *Circulation* 2005; 112: 2735–2752.
36. Tondo L, Isacsson G, Baldessarini R. Suicidal behaviour in bipolar disorder: risk and prevention. *CNS Drugs* 2003; 17(7):491-511. <https://doi-org.bibliopass.unito.it/10.2165/00023210-200317070-00003>
37. Manchia M, Hajek T, O'Donovan C, Deiana V, Chillotti C, Ruzickova M, Del Zompo M, Alda M. Genetic risk of suicidal behavior in bipolar spectrum disorder: analysis of 737 pedigrees. *Bipolar Disord* 2013; 15: 496–506. <https://doi-org.bibliopass.unito.it/10.1111/bdi.12088>
38. Tidemalm D, Haglund A, Karanti A, Landén M, Runeson B. Attempted Suicide in Bipolar Disorder: Risk Factors in a Cohort of 6086 Patients. *PLoS One* 2014; 9(4):e94097. <https://doi-org.bibliopass.unito.it/10.1371/journal.pone.0094097>
39. Vuksan-Ćusa B, Marčinko D, Nađ S, Jakovljević M. Differences in cholesterol and metabolic syndrome between bipolar disorder men with and without suicide attempts. *Prog Neuropsychopharmacol Biol Psychiatry* 2009; 33(1):109-12. <https://doi-org.bibliopass.unito.it/10.1016/j.pnpbp.2008.10.017>
40. Yen CF, Cheng CP, Ko CH, Yen JY, Huang CF, Chen CS. Suicidality and its association with insight and neurocognition in taiwanese patients with bipolar I disorder in remission. *J. Nerv Ment Dis* 2008; 196(6):462-7. <https://doi-org.bibliopass.unito.it/10.1097/NMD.0b013e3181775a3d>
41. De Abreu LN, Nery FG, Harkavy-Friedman JM, De Almeida KM, Gomes BC, Oquendo MA. Suicide attempts are associated with worse quality of life in patients with bipolar disorder type I. *Compr Psychiatry* 2012; 53(2):125-9. <https://doi.org/10.1016/j.comppsy.2011.03.003>.
42. Galvez JF, Bauer IE, Sanches M, Wu HE, Hamilton JE, Mwangi B, Kapczinski FP, Zunta-Soares G, Soares JC. Shared clinical associations between obesity and impulsivity in rapid cycling bipolar disorder: A systematic review. *JAD* 2012; 168 (2014) 306–313. <https://doi-org.bibliopass.unito.it/10.1016/j.jad.2014.05.054>

Table 1 Socio-demographic and clinical characteristics of the total sample (n = 842) and differences in socio-demographic and clinical characteristics between suicide attempters (n = 202) and non attempters (n = 640).

Characteristics	Total sample (n = 842)	Suicide attempters (n = 202)	Non-suicide attempters (n = 640)	F/ χ^2	df	p
Age (years), mean \pm <i>sd</i>	49.9 \pm 15.7	50.8 \pm 14.7	49.6 \pm 16.0	1.017	1	0.313
Education (years), mean \pm <i>sd</i>	11.9 \pm 4.2	11.8 \pm 4.1	11.9 \pm 4.2	0.184	1	0.668
Sex, n (%)						
Male	348 (41.3)	77 (38.1)	271 (42.3)	1.130	1	0.288
Female	494 (58.7)	125 (61.9)	369 (57.7)			
Bipolar disorder, type, n (%)						
Bipolar I	352 (41.8)	73 (36.1)	279 (43.6)	5.072	4	0.280
Bipolar II	459 (54.5)	123 (60.9)	336 (52.5)			
Bipolar NOS	28 (3.3)	6 (3.0)	22 (3.4)			
Age of onset (years), mean \pm <i>sd</i>	29.3 \pm 12.4	28.4 \pm 11.8	29.6 \pm 12.6	1.528	1	0.217
Duration of illness (years), mean \pm <i>sd</i>	20.5 \pm 14.2	22.4 \pm 14.1	19.9 \pm 14.2	4.858	1	0.028
Duration of untreated illness (years), mean \pm <i>sd</i>	13.2 \pm 12.5	14.3 \pm 13.0	12.9 \pm 12.4	1.804	1	0.180
Manic episodes (number), mean \pm <i>sd</i>	1.3 \pm 2.5	1.0 \pm 2.1	1.4 \pm 2.6	3.154	1	0.077
Hypomanic episodes (number), mean \pm <i>sd</i>	2.6 \pm 3.4	3.3 \pm 4.3	2.3 \pm 3.1	11.965	1	0.001
Depressive episodes (number), mean \pm <i>sd</i>	5.0 \pm 4.2	6.0 \pm 4.4	4.7 \pm 4.1	13.741	1	<0.001
Lifetime psychiatric comorbidities, n (%)	365 (43.3)	101 (50.0)	264 (41.3)	4.787	1	0.029
Family history of mood disorders, n (%)	483 (57.6)	119 (59.2)	364 (57.1)	0.266	1	0.606
Family history of suicide, n (%)	11 (6.5)	3 (6.4)	8 (7.7)	0.080	1	0.774
Lifetime medical comorbidity, n (%)	402 (51.1)	112 (58.0)	290 (48.9)	4.855	1	0.028
Weight (kg), mean \pm <i>sd</i>	72.6 \pm 16.0	72.6 \pm 16.6	72.6 \pm 15.8	0.001	1	0.974
BMI (kg/m ²), mean \pm <i>sd</i>	28.5 \pm 5.8	25.8 \pm 5.1	29.5 \pm 5.2	0.386	1	0.535
Waist circumference (cm), mean \pm <i>sd</i>	94.2 \pm 16.7	94.6 \pm 16.1	94.1 \pm 16.9	0.071	1	0.790
Serum lipid levels (mg/dl), mean \pm <i>sd</i>						
Cholesterol	195.2 \pm 44.7	193.5 \pm 46.1	195.8 \pm 44.2	0.290	1	0.591
Tryglicerides	134.2 \pm 77.5	135.3 \pm 85.4	133.8 \pm 74.7	0.046	1	0.829
HDL cholesterol	51.9 \pm 16.5	51.2 \pm 18.1	52.1 \pm 15.9	0.358	1	0.550
Glycemia (mg/dl), mean \pm <i>sd</i>	86.4 \pm 22.8	88.1 \pm 23.0	85.9 \pm 22.8	1.214	1	0.271
Systolic arterial pressure (mmHg), mean \pm <i>sd</i>	122.9 \pm 12.5	122.1 \pm 12.5	123.2 \pm 12.5	1.006	1	0.316
Diastolic arterial pressure (mmHg), mean \pm <i>sd</i>	79.1 \pm 8.8	77.9 \pm 8.7	79.4 \pm 8.8	3.794	1	0.052
Metabolic syndrome, n (%)	198 (29.3)	58 (33.1)	140 (28.0)	1.654	1	0.198
Abdominal obesity, n (%)	259 (49.4)	73 (51.4)	186 (48.7)	0.306	1	0.580
Low HDL cholesterol, n (%)	256 (39.1)	78 (46.2)	178 (36.7)	4.701	1	0.030
Elevated blood pressure, n (%)	321 (47.9)	85 (48.6)	236 (47.7)	0.387	1	0.824
Impaired fasting glucose, n (%)	117 (17.5)	37 (21.5)	80 (16.1)	2.876	1	0.237
Hypertriglyceridemia, n (%)	194 (29.1)	46 (26.7)	148 (29.9)	1.361	1	0.715

NOS: Not Otherwise Specified

BMI: Body Mass Index

Table 2 Differences in socio-demographic and clinical characteristics between violent (n = 52) and nonviolent (n = 118) suicide attempters.

Characteristics	Violent attempt (n = 52)	Nonviolent attempt (n = 118)	F/ χ^2	df	p
Age (years), mean \pm sd	49.9 \pm 15.9	50.2 \pm 14.1	0.062	1	0.831
Education (years), mean \pm sd	11.8 \pm 3.8	11.5 \pm 4.1	0.179	1	0.673
Sex, n (%)					
Male	34 (65.4)	33 (28.0)	21.16	1	<0.001*
Female	18 (34.6)	85 (72.0)			
Bipolar disorder, type, n (%)					
Bipolar I	17 (32.7)	44 (37.3)	0.69	2	0.707
Bipolar II	34 (65.4)	70 (59.3)			
Bipolar NOS	1 (1.9)	4 (3.4)			
Age of onset (years), mean \pm sd	28.7 \pm 13.7	27.9 \pm 11.5	0.138	1	0.711
Duration of illness (years), mean \pm sd	21.3 \pm 14.3	22.4 \pm 13.6	0.222	1	0.638
Duration of untreated illness (years), mean \pm sd	14.7 \pm 14.3	14.7 \pm 12.9	0.001	1	0.976
Manic episodes (number), mean \pm sd	0.8 \pm 1.5	1.2 \pm 2.3	0.839	1	0.361
Hypomanic episodes (number), mean \pm sd	3.1 \pm 3.8	3.1 \pm 4.5	0.001	1	0.994
Depressive episodes (number), mean \pm sd	5.7 \pm 4.0	6.2 \pm 4.8	0.460	1	0.499
Lifetime psychiatric comorbidity, n (%)	23 (44.2)	56 (47.4)	0.151	1	0.697
Family history of mood disorders, n (%)	28 (53.8)	71 (60.7)	0.69	1	0.405
Family history of suicide, n (%)	3 (6.4)	8 (7.7)	0.080	1	0.774
Current psychopharmacological treatment, n (%)					
Monotherapy	6 (11.5)	21 (17.8)	3.240	2	0.198
Combination therapy	45 (88.5)	97 (82.2)			
None	1 (1.9)	0 (0)			
Lithium, n (%)	31 (59.6)	76 (65.0)	0.442	1	0.506
Anticonvulsivants, n (%)	27 (51.9)	53 (44.9)	0.771	1	0.399
First-generation antipsychotics, n (%)	8 (15.4)	7 (5.9)	4.009	1	0.074
Second-generation antipsychotics, n (%)	30 (57.7)	60 (50.8)	0.679	1	0.410
Antidepressants, n (%)	19 (36.5)	48 (40.7)	0.259	1	0.611
Lifetime medical comorbidity, n (%)	32 (61.5)	70 (59.3)	0.044	1	0.083
Weight (kg), mean \pm sd	80.5 \pm 18.3	69.4 \pm 14.7	16.6	1	<0.001*
BMI (kg/m ²), mean \pm sd	27.8 \pm 5.6	25.2 \pm 4.7	9.34	1	0.003*
Waist circumference (cm), mean \pm sd	98.7 \pm 18.5	92.4 \pm 14.3	0.968	1	0.032*
Serum lipid levels (mg/dl), mean \pm sd					
Cholesterol	190.6 \pm 49.2	194.6 \pm 45.0	0.234	1	0.629
Tryglicerides	154.2 \pm 95.1	131.6 \pm 83.2	2.272	1	0.134
HDL cholesterol	47.6 \pm 15.4	52.0 \pm 18.9	1.955	1	0.164
Glycemia (mg/dl), mean \pm sd	89.1 \pm 23.8	86.9 \pm 23.3	0.296	1	0.587
Systolic arterial pressure (mmHg), mean \pm sd	124.1 \pm 11.9	120.8 \pm 12.4	2.529	1	0.114
Diastolic arterial pressure (mmHg), mean \pm sd	78.9 \pm 8.8	77.5 \pm 8.8	0.958	1	0.327
Metabolic syndrome, n (%)	20 (35.7)	36 (41.3)	1.13	1	0.288
Abdominal obesity, n (%)	21 (40.4)	49 (41.5)	1.704	1	0.300
Low HDL cholesterol, n (%)	19 (36.5)	54 (45.8)	0.605	1	0.437
Elevated blood pressure, n (%)	27 (51.9)	50 (42.4)	1.006	1	0.316
Impaired fasting glucose, n (%)	12 (23.1)	20 (16.9)	0.789	1	0.374
Hypertriglyceridemia, n (%)	13 (25.0)	31 (26.3)	0.124	1	0.724

* remained significant after Bonferroni correction: p<0.0017

NOS: Not Otherwise Specified

BMI: Body mass index

Table 3 Relationship between potential explanatory variables and violent suicide methods: results from the logistic regression analysis

Dependent variables	B	S.E.	Wald	ORa	95% CI OR	p
Gender (Male)	1.820	0.545	11.147	6.174	(2.121-17.972)	0.001
Body weight	-0.002	0.033	0.004	0.998	(0.935-1.065)	0.948
Body Mass Index	0.086	0.096	0.807	1.090	(0.903-1.315)	0.369
Abdominal circumference	-0.037	0.018	4.309	0.964	(0.931-0.998)	0.036